

Aireworth Vets 
SMALL ANIMAL  

CANINE BLADDER STONES



Aireworth Vets 
SMALL ANIMAL  

If you have any further queries regarding the welfare of your pet, please do not hesitate to ask a member of staff

We are happy to assist!

Aireworth Road, Keighley, West Yorkshire BD21 4DJ
Tel: 01535 602988
Fax: 01535 691498
Email: Admin@aireworthvets.co.uk

Aireworth Veterinary Centre
Aireworth Road
Keighley
BD21 4DJ

Tel: 01535 602988
Fax: 01535 691498

Email: admin@aireworthvets.co.uk

What are bladder stones?

Bladder stones, more correctly called uroliths, or cystic calculi are rock-like collections of minerals that form in the urinary bladder. They may occur as a large, single stone or as collections of stones the size of large grains of sand or gravel.

Are these the same as gall stones or kidney stones?

No. Gall stones are in the gall bladder, and kidney stones are in the kidney. Although the kidneys and urinary bladder are both part of the urinary system, kidney stones are usually unrelated to bladder stones.

What problems do bladder stones cause?

The two most common signs of bladder stones are haematuria (blood in the urine) and dysuria (straining to urinate). Haematuria occurs because the stones mechanically irritate the bladder wall, causing bleeding from its surface. Dysuria occurs when stones obstruct the passage of urine out of the bladder. Large stones may cause a partial obstruction at the point where the urine leaves the bladder and enters the urethra; small stones may flow with the urine into the urethra and cause an obstruction there.

When an obstruction occurs, the bladder cannot be emptied and this is very painful. Your dog may cry in pain, especially if pressure is applied to the abdominal wall. Haematuria and dysuria are the most common signs seen in dogs with bladder stones but with obstruction there is usually pain as well. We know this because when bladder stones are removed surgically, many owners tell us how much better and more active their dog feels.

Why do they form?

There are several theories to explain the formation of bladder stones. Each is feasible in some circumstances, but there is probably an interaction of more than one of them in each dog. The most commonly accepted theory is called the Precipitation-Crystallisation Theory. This theory states that one or more stone-forming crystalline compounds is present in elevated levels in the urine. This may be due to abnormalities in diet or due to some previous disease in the bladder, especially infection with bacteria. Sometimes the condition may be due to a fault in body chemistry. When the amount of this compound reaches a threshold level, the urine is said to be supersaturated. This means that the level of the compound is so great that it cannot all be dissolved in the urine, so it precipitates and forms tiny crystals. These crystals stick together, usually due to mucus-like material within the bladder, and stones gradually form. As time passes, the stones enlarge and increase in number.

How are they diagnosed?

Most dogs that have bladder infections do not have bladder stones. These dogs will often have blood in the urine and will strain to urinate, the same symptoms as a dog with bladder stones. Therefore, we do not suspect bladder stones just based on these clinical signs.

Some bladder stones can be palpated (felt with the fingers) through the abdominal wall. However, failure to palpate them does not rule them out. Most bladder stones are visible on radiographs (x-rays) or an ultrasound examination. These procedures are performed if stones are suspected. This includes dogs that show unusual pain when the bladder is palpated, dogs that have recurrent haematuria and dysuria, or dogs that have recurrent bacterial infections in the bladder.


Some bladder stones are not visible on radiographs. They are said to be radiolucent. This means that their mineral composition is such that they do not reflect the x-ray beam. These stones may be found with an ultrasound examination or with special radiographs that are made after placing a special dye (contrast material) in the bladder.

How are bladder stones treated?

The fastest way is to remove them surgically. This requires major surgery in which the abdomen and bladder are opened. Following two to four days of recovery, the dog is relieved of pain and dysuria. The haematuria will often persist for a few more days, then it stops. Surgery is not the best option for all patients; however, those with urethral obstruction and those with bacterial infections associated with the stones should be operated upon unless there are other health conditions that prohibit surgery.

Can bladder stones be prevented?

The answer is a qualified "yes." There are at least four types of bladder stones, based on their chemical composition. If stones are removed surgically or if some small ones pass in the urine, they should be analysed for their chemical composition. This will permit us to determine if a special diet will be helpful in preventing recurrence. If a bacterial infection causes stone formation, it is recommended that periodic urinalyses and urine cultures be performed to determine when antibiotics should be given.



Canine bladder stones



If you have any queries please contact us.